

ON THE PROGENITOR OF SUPERNOVA 2001DU IN NGC 1365¹

SCHUYLER D. VAN DYK

IPAC/Caltech, Mailcode 100-22, Pasadena CA 91125

WEIDONG LI AND ALEXEI V. FILIPPENKO

Department of Astronomy, 601 Campbell Hall, University of California, Berkeley, CA 94720-3411

To appear in PASP (2003 April).

ABSTRACT

Hubble Space Telescope (HST) WFPC2 archival images obtained years prior to the explosion of the Type II-plateau supernova (SN) 2001du in NGC 1365 previously have been analyzed to isolate the progenitor star. The SN site was located using precise astrometry applied to the *HST* images, with significant uncertainty, leaving three possible progenitor candidates. Images of the fading SN have recently become publicly available in the *HST* archive, allowing us to pinpoint the SN's exact location on the pre-explosion images. We show that the SN occurred in very close proximity to one of the blue candidate stars, but we argue that this star is not the actual progenitor. Instead, the progenitor was not detected on the pre-SN images, and we constrain the progenitor's mass to be less than $13_{-4}^{+7} M_{\odot}$. This is consistent with previous constraints on the progenitor masses of other Type II-plateau supernovae (SNe), suggesting that such SNe arise from the iron core collapse of massive stars at the lower extreme of the possible mass range.

Subject headings: supernovae: general — supernovae: individual (SN 2001du) — stars: massive — stars: evolution — stars: variables: other — galaxies: individual (NGC 1365)

1. INTRODUCTION

Supernova 2001du was visually discovered by R. Evans (2001) on August 24.7 (UT dates are used throughout this paper) at the western end of the bar in the spiral galaxy NGC 1365 (about 90'' W and 10'' S of the nucleus). Smartt, Kilkenny, & Meikle (2001) classified the SN as Type II; Wang et al. (2001), from a spectrum obtained on August 30 UT, confirmed the SN as Type II-plateau (II-P) before optical maximum. NGC 1365 has also been host to SN 1957C (of unknown type) and the Type Ic SN 1983V. Using *Hubble Space Telescope (HST)* observations of Cepheids, Silbermann et al. (1999) determined the distance modulus of NGC 1365 to be $\mu = 31.3 \pm 0.2$ mag, corresponding to a distance of about 18.2 Mpc.

Van Dyk, Li, & Filippenko (2003) recently measured the absolute position of SN 2001du and, applying an independent astrometric grid to pre-SN *HST* archival WFPC2 images, isolated three possible candidate progenitor stars within the uncertainty of the measured SN position: one reddish star with a detection in both the F555W and F814W bands, and two stars with only F555W detections, implying bluer colors for these latter two stars. Also, multi-band WFPC2 images of the SN were obtained for *HST* program GO-9041 (PI: S. Smartt) in 2001 November. These images are now in the public domain, and in this paper we analyze them and compare them to the pre-SN images to further constrain the nature of the SN progenitor. (Smartt et al. 2003 have also recently analyzed their images and, in a somewhat similar manner, come to the same basic conclusion that we reach here.)

2. ANALYSIS

Pre-SN images, each of total exposure time 300 s in the bands F336W, F555W, and F814W, were obtained by GTO-5222 on 1995 January 15. Late-time multi-band images of the SN were obtained by GO-9041 on 2001 November 26. These include deep images in F336W, F439W, F555W, F675W, and F814W, as well as shorter exposures in F336W, F555W, and F814W. The available *HST* data are summarized in Table 1, with total exposure times given for each band.

In Van Dyk et al. (2003) we adopted the position of the SN, measured from an image obtained with the YALO 1-m telescope at CTIO (and available on the Internet), as $\alpha = 3^h 33^m 29^s.15$, $\delta = -36^\circ 08' 31''.5$ (J2000), with total uncertainty $\pm 0''.9$. The SN site is located on the WF3 chip. In Figure 1 we reproduce Figure 21 from Van Dyk et al., showing the SN environment in the F555W band. In Figure 2 we show the late-time F555W image (the SN is nearly centered on the PC chip for all the GO-9041 exposures). As one can see, the actual SN position is within the $0''.9$ error circle, which provides additional confidence in the astrometric method used by Van Dyk et al. when isolating the various SN sites².

We initially used the routine HSTphot (Dolphin 2000a,b) to obtain photometry of the objects imaged on the PC chip. HSTphot automatically accounts for WFPC2 point-spread function (PSF) variations and charge-transfer effects across the chips, zeropoints, aperture corrections, etc., and can return as output magni-

¹BASED ON OBSERVATIONS MADE WITH THE NASA/ESA *HUBBLE SPACE TELESCOPE*, OBTAINED FROM THE DATA ARCHIVE OF THE SPACE TELESCOPE SCIENCE INSTITUTE, WHICH IS OPERATED BY THE ASSOCIATION OF UNIVERSITIES FOR RESEARCH IN ASTRONOMY, INC., UNDER NASA CONTRACT NAS 5-26555.

²Applying the same astrometric method used by Van Dyk et al. (2003) to the mosaic of the deep F555W exposure, we measure a position for SN 2001du to be $\alpha = 3^h 33^m 29^s.14$, $\delta = -36^\circ 08' 31''.3$ (J2000), $\pm 0''.2$ in each coordinate.

tudes transformed from the *HST* flight system to standard Johnson-Cousins bands. Van Dyk et al. (2003) discuss the similarity in the results from HSTphot, compared with those from the packages DoPHOT (Schechter, Mateo, & Saha 1993) and DAOPHOT/ALLSTAR (Stetson 1987, 1992).³

The SN profile is saturated in the longer F555W (*V*), F675W (*R*), and F814W (*I*) exposures. Thus, no brightness measurement is possible. The SN is not saturated in the shorter F555W and F814W exposures, the long F336W (*U*) and F439W (*B*) exposures, and also the short F336W exposure. However, the results from HSTphot for the F336W exposures varied, depending on how these exposures were combined with the other bands as input into the routine, and the results from the long and short exposures for this band were also discrepant with each other (the results ranged from about $m_{F336W} = 18.9$ to 18.7 mag for different runs of HSTphot). We do not know why this occurred, but we speculate that it possibly may be due to the low signal-to-noise ratio for both exposure sets. For F336W we then applied to both the long and short exposures DAOPHOT/ALLSTAR within IRAF⁴ as a check, using a TinyTim PSF (Krist 1995), the zeropoint from Holtzman et al. (1995), and aperture corrections (~ 0.00 mag) derived from artificial star tests. (Checks for the F439W and short F555W and F814W exposures with DAOPHOT show good agreement with the HSTphot results.)

We obtained $m_{F336W} = 18.96 \pm 0.10$ and 18.87 ± 0.04 mag for the short and long exposures, respectively. We then averaged these DAOPHOT magnitudes with the HSTphot magnitudes at F336W, and obtained $m_{F336W} = 18.86$ mag. The SN was quite red by late November 2001, with $m_{F336W} - m_{F439W} = 1.82$ mag. Whether using the transformations from flight system to Johnson-Cousins magnitudes from Holtzman et al. (1995), Dolphin (2000b), or via SYNPHOT applied to the Bruzual Spectral Synthetic Atlas (see Van Dyk, Filippenko, & Li 2002), this color is too red to obtain a reliable transformation from m_{F336W} to *U*. However, we do apply a transformation, with uncertainty ~ 0.2 mag. The transformations for all other bands are from Dolphin (2000b), applied internally within HSTphot. The final results for the photometry of SN 2001du in the *HST* images are presented in Table 2.

In Figure 3 we show preliminary, unpublished *BVI* light curves for SN 2001du, based on data from the YALO 1-m telescope (N. Suntzeff & K. Krisciunas, 2003, private communication), and we include the *UBVI* photometry from the *HST* data. For comparison we show the *B* and *V* light curves for the plateau-type SN (II-P) 1999em in NGC 1637 (Leonard et al. 2002a). It is clearly evident that SN 2001du is indeed a SN II-P as well. If this SN were a linear-type event (II-L), it should have markedly declined in brightness by the time of the *HST* observations (~ 94 days since discovery).

3. DISCUSSION

³In fact, here we update the private communication by D. C. Leonard 2002, regarding the very good agreement between the results of HSTphot photometry on images of NGC 3351, compared with those obtained by Graham et al. (1997) using DAOPHOT; Leonard now finds $\delta V = 0.015 \pm 0.022$ and $\delta I = 0.036 \pm 0.018$ mag.

⁴IRAF (Image Reduction and Analysis Facility) is distributed by the National Optical Astronomy Observatories, which are operated by the Association of Universities for Research in Astronomy, Inc., under cooperative agreement with the National Science Foundation.

⁵NED is the NASA/IPAC Extragalactic Database, <http://nedwww.ipac.caltech.edu>.

In Van Dyk et al. (2003) we had isolated three faint objects, A–C, detected and determined to be stellar by HSTphot, within the error circle around the SN position. Stars A and B, to the south and west, respectively, are both blue, with $m_{F555W} (\sim V) = 24.30 \pm 0.21$ mag and 25.02 ± 0.32 mag, respectively, and no detections at F814W ($\sim I$). (Blue, here, is relative, and includes stars with roughly early-G spectral type or earlier, based on the detection limits; see below.) Star C, to the east, is relatively red, with $V = 24.44 \pm 0.23$ and $V - I = 1.03 \pm 0.30$ mag. We had assumed that, since SN 2001du is of Type II-P, the most plausible of the three candidates is the redder star. However, that was argued to be still unlikely, based on the assumed luminosity and color of Star C.

We have used the late-time SN images, specifically the *V* image, and attempted to match them up with the pre-SN images. This was done by using the task *geotran* in IRAF to transform both the late-time SN images to the pre-SN images and vice versa. We can therefore locate the position of the SN on the pre-SN images as being 0.70 ± 0.15 WF pixel ($0''.07$) northeast of the position of Star B. It would be tempting to assign Star B, then, as the progenitor of SN 2001du. However, (1) the offset of the SN position from Star B (which corresponds spatially to 6.2 pc at the distance of NGC 1365) is significantly large, and the accuracy of the measurement is appreciably high; (2) SNe II-P are thought to arise from the explosions of red supergiants; and (3) no evidence exists that SN 2001du is peculiar in any way and that it arose from a bluer, more compact star like the progenitor of SN 1987A (E. Cappellaro, 2002, private communication). Thus, it is more likely, as we previously had suspected (Van Dyk et al. 2003), that the progenitor simply is not detected in the pre-SN images.

We can use the detection limits of both the F555W and F814W pre-SN images to constrain the nature of the progenitor. We do this by inferring limits on the luminosity and, therefore, the mass from the limits on the star's apparent magnitude. These HSTphot detection limits in the SN environment are $m_{F555W} > 25.19$ and $m_{F814W} > 24.25$ mag. Before converting into absolute magnitude, we must first assess the amount of extinction to the SN. In Van Dyk et al. (2003) we attempted to use stars in the environment to estimate the reddening. The Galactic extinction toward NGC 1365 is $A_V = 0.07$ mag (Schlegel, Finkbeiner, & Davis 1998; and NED⁵). We had found that stars in the larger SN environment implied larger amounts of extinction, $A_V \approx 2.5$ mag. (Unfortunately, although the post-SN images are deeper than the pre-SN ones, the SN is far too bright, and its image is spread over too many surrounding pixels, to perform a more thorough study of the environment.)

Using the light curve in Figure 3 we can actually derive more specific information about the possible reddening toward the SN from the SN itself (which is information that was not available to us previously). The *HST* photometry of the SN on November 26 is about 94 days after discovery, i.e., near the end of the light-curve plateau, where we

can still estimate the SN's color temperature with reasonable accuracy (e.g., Hamuy et al. 2001). These colors are $B - V = 1.42$ and $V - I = 0.91$ mag. For comparison, the colors of the very well-studied Type II-P SN 1999em at the same age (about 95 days after discovery) are $B - V = 1.47$ and $V - I = 0.89$ mag (Leonard et al. 2002a). Leonard et al. and Hamuy et al. (2001) conclude that SN 1999em is reddened by $E(B - V) = 0.10$ mag. Since $B - V$ is affected by line blanketing in the B band (e.g., Hamuy et al. 2001), the small difference in $B - V$ (SN 1999em is 0.05 mag redder than SN 2001du) could be due to differences in metallicity between the two SNe, which might affect this color. The marginally redder $V - I$ color of SN 2001du (by 0.02 mag) may imply that this SN is slightly more reddened than SN 1999em. However, we consider these differences in color between the two SNe to be small and likely within the uncertainties in the colors, so we adopt $E(B - V) \approx 0.1$ mag for SN 2001du, as found for SN 1999em.

If we assume the Cardelli, Clayton, & Mathis (1989) reddening law, then $A_V \approx 0.3$ mag and $A_I \approx 0.2$ mag for SN 2001du. For the distance modulus $\mu = 31.3$ mag, $M_V > -6.4$ and $M_I > -7.0$ mag for the progenitor star. In a similar vein to the estimation of the progenitor mass limit for SN II-P 1999em by Smartt et al. (2002) and SN II-P 1999gi by Smartt et al. (2001), the absolute magnitude limits can be converted to the luminosity of the likely supergiant progenitor, assuming the full range of possible stellar surface temperatures. This is accomplished using a similar relation to equation (1) in Smartt et al. (2001, 2002), where we also have assumed the temperatures and stellar bolometric corrections for supergiants from Drilling & Landolt (2000) and have computed, using SYNPHOT within IRAF and the Bruzual Spectral Atlas, the conversions of m_{F555W} and m_{F814W} to Johnson V for the range in spectral type. The luminosity limits for both F555W (V) and F814W (I) are shown in Figure 4. Stars with luminosities brighter than these limits (shown by the *heavy lines*) should have been detected in the pre-SN images.

For comparison we have plotted (also in a similar fashion to Smartt et al.) the model stellar evolutionary tracks for a range of masses from Lejeune & Schaerer (2001), assuming enhanced mass loss for the most massive stars and that a metallicity $Z = 0.04$ is appropriate for the SN environment. Indication for a metallicity somewhat greater than solar in this sector of NGC 1365 comes from the abundance study by Roy & Walsh (1997) of H II regions in the galaxy. The closest H II regions to the SN site are their Nos. 20 and 21, where at this radial distance from the nucleus, the O/H abundance is ~ 9 dex or, in fact, slightly greater, based on model assumptions (the solar O/H abundance is 8.8 dex; Grevesse & Sauval 1998).

The luminosity limits at each band in Figure 4 show that the pre-SN images are not sensitive to the bluest high-mass supergiants, so we cannot be absolutely sure that such a star is not the progenitor. If we exclude this possibility, the V band sets an upper limit to the progenitor mass of $\sim 13 M_\odot$, although this ranges upward to as high as $\sim 20 M_\odot$, depending on the effective temperature. The I band is more sensitive to less-massive red supergiants, ranging downward to $\sim 9 M_\odot$ (possibly even less) at the lowest effective temperatures. These results from Figure 4

are roughly consistent with what we showed in Figure 22 of Van Dyk et al. (2003). We therefore estimate that the upper mass limit for the SN 2001du progenitor is $13^{+7}_{-4} M_\odot$.

This limit is consistent with the limit Leonard et al. (2002b) find for SN II-P 1999gi ($15^{+5}_{-3} M_\odot$) and the $\sim 17 M_\odot$ limit one finds for SN II-P 1999em, if the distance to the SN 1999em host is assumed to be the Cepheid distance, 11.5 ± 0.9 Mpc (Leonard et al. 2003). Low masses for the progenitors of SNe II-P, relative to that indicated for SNe II-L (linear) and SNe IIn (narrow), are also inferred from their optical properties (e.g., Schlegel 1996; Filippenko 1997; Chugai & Utrobin 2000) and suggested by the inferred paucity of circumstellar matter from radio and X-ray observations (e.g., Weiler et al. 1989; Schlegel 2001; Pooley et al. 2002).

4. CONCLUSIONS

Comparing *HST* archival WFPC2 images obtained of the Type II-P SN 2001du in NGC 1365 at late times with images obtained before explosion, we have located the position of the SN progenitor star to within 1 WF chip pixel of the candidate Star B identified by Van Dyk et al. (2003). This star, however, is more than likely *not* the progenitor. We use the detection limits of the pre-SN F555W and F814W images to place constraints on the progenitor's nature; specifically, $M \lesssim 13^{+7}_{-4} M_\odot$. (Smartt et al. 2003 find a similar mass limit of $M = 15 M_\odot$.) This limit is consistent with what has been found and inferred for other SNe II-P.

The proximity of the SN to the blue Star B implies that the progenitor may have been part of a small stellar association (the separation between the progenitor and Star B is likely too large for the blue star to be a binary companion). The faint emission underlying Stars A, B, and C identified by Van Dyk et al. (2003) suggests that an association complex may lie in the SN environment. It would be interesting to image the SN again in multiple bands with *HST* at high spatial resolution, presumably with the Advanced Camera for Surveys (ACS), when the SN has significantly dimmed. The stars in the environment of the fading SN would then become more obvious, and the age and mass of the progenitor star might be further constrained, based on the characteristics of its surviving neighbors.

This research made use of the NASA/IPAC Extragalactic Database (NED) which is operated by the Jet Propulsion Laboratory, California Institute of Technology, under contract with NASA, and the LEDA database (<http://leda.univ-lyon.fr>). The work of A.V.F.'s group at UC Berkeley is supported by NASA grants AR-8754, AR-9529, and GO-8602 from the Space Telescope Science Institute, which is operated by AURA, Inc., under NASA contract NAS5-26555. We thank D. C. Leonard and M. Hamuy for useful discussions. We also thank the referee, K. Krisciunas, for helpful comments and for providing unpublished ground-based photometry of SN 2001du.

REFERENCES

- Cardelli, J. A., Clayton, G. C., & Mathis, J. S. 1989, *ApJ*, 345, 245
 Chugai, N. N., & Utrobin, V. P. 2000, *A&A*, 354, 557
 Dolphin, A. E. 2000a, *PASP*, 112, 1383
 Dolphin, A. E. 2000b, *PASP*, 112, 1397
 Drilling, J.S., & Landolt, A.U. 2000, in *Allen's Astrophysical Quantities*, 4th ed., ed. A. N. Cox (New York: AIP), 381
 Evans, R. 2001, *IAU Circ.* 7690
 Filippenko, A. V. 1997, *ARA&A*, 35, 309
 Graham, J. A., et al. 1997, *ApJ*, 477, 535
 Grevesse, N., & Sauval, A.J. 1998, *Space Sci. Rev.*, 85, 161
 Hamuy, M., et al. 2001, *ApJ*, 558, 615
 Holtzman, J. A., Burrows, C. J., Casertano, S., Hester, J. J., Trauger, J. T., Watson, A. M., & Worthey, G. 1995, *PASP*, 107, 1065
 Krist, J. 1995, in *Calibrating Hubble Space Telescope: Post Servicing Mission* (Baltimore: STScI), 311
 Lejeune, T., & Schaerer, D. 2001, *A&A*, 366, 538
 Leonard, D. C., et al. 2002a, *PASP*, 114, 35
 Leonard, D. C., Kanbur, S. M., Ngeow, C. C., & Tanvir, N. R. 2003, *ApJ*, submitted
 Leonard, D. C., et al. 2002b, *AJ*, 124, 2490
 Pooley, D., et al. 2002, *ApJ*, 572, 932
 Roy, J.-R., & Walsh, J. R. 1997, *MNRAS*, 288, 715
 Schechter, P. L., Mateo, M., & Saha, A. 1993, *PASP*, 105, 1342
 Schlegel, E. M. 1996, *AJ*, 111, 1660
 Schlegel, E. M. 2001, *ApJ*, 556, L29
 Schlegel, D. J., Finkbeiner, D. P., & Davis, M. 1998, *ApJ*, 500, 525
 Silbermann, N. A., et al. 1999, *ApJ*, 515, 1
 Smartt, S. J., Kilkenny, D., & Meikle, P. 2001, *IAUC* 7704
 Smartt, S. J., Gilmore, G. F., Trentham, N., Tout, C. A., & Frayn, C. M. 2001, *ApJ*, 556, L29
 Smartt, S. J., Gilmore, G. F., Tout, C. A., & Hodgkin, S. T. 2002, *ApJ*, 565, 1089
 Smartt, S. J., Maund, J. R., Gilmore, G. F., Tout, C. A., Kilkenny, D., & Benetti, S. 2003, *MNRAS*, submitted
 Stetson, P. B. 1987, *PASP*, 99, 191
 Stetson, P. B. 1992, in *ADASS (ASP Conf. Ser. 25)*, ed. D. M. Worral, C. Bimesderfer, & J. Barnes (San Francisco: ASP), 297
 Van Dyk, S. D., Filippenko, A. V., & Li, W. D. 2002, *PASP*, 114, 701
 Van Dyk, S. D., Li, W. D., & Filippenko, A. V., 2003, *PASP*, 115, 1
 Wang, L., Baade, D., Fransson, C., Höflich, P., Lundqvist, P., & Wheeler, J. C. 2001, *IAU Circ.* 7704
 Weiler, K. W., Panagia, N., Sramek, R. A., van der Hulst, J. M., Roberts, M. S., & Nguyen, L. 1989, *ApJ*, 336, 421

TABLE 1
SUMMARY OF *HST* OBSERVATIONS

Date (UT)	Filter	Exp. Time (s)	<i>HST</i> Program	SN Saturated?
1995 Jan 15	F336W	300 ^a	GTO-5222	...
	F555W	300		...
	F814W	300		...
2001 Nov 26	F336W	100	GO-9041	N
		460		N
	F439W	400		N
	F555W	40		N
		600		Y
	F675W	400		Y
	F814W	40		N
		600		Y

^aThe exposure times for all three bands in this 1995 Jan 15 dataset were erroneously listed in Van Dyk et al. (2003) as 100 s. The actual total exposure time of 300 s results from the combination of a trio of 100-s exposures. Note that HSTphot automatically reads the correct total exposure time from the images during processing, so the reported results are unaffected by this error.

TABLE 2
PHOTOMETRY OF THE SN 2001DU *HST* IMAGES

U [m_{F336W}]	B [m_{F439W}]	V [m_{F555W}]	R [m_{F675W}]	I [m_{F814W}]
18.6(0.2) ^a [18.86] ^b	16.72(0.06) [17.04]	15.30(0.00) [15.32]		14.39(0.00) [14.43]

^aThe value in parentheses for each entry in this row is the uncertainty in the transformed Johnson-Cousins magnitude.

^bThe value in brackets for each entry in this row is the adopted flight-system magnitude, derived from HSTphot, DAOPHOT, or both.

Fig. 1.— The site of SN 2001du in the archival 300-s F555W image from 1995 January 15. The error circle has radius $0''.9$. The three stars, A–C, within the circle are indicated with tickmarks. This is a reproduction of Fig. 21 from Van Dyk et al. (2003).

Fig. 2.— SN 2001du, as seen in the 600-s F555W image from 2001 November 26, shown to the same scale and orientation as in Figure 1. The error circle from Figure 1 is also shown. The supernova is saturated in this image.




Fig. 3.— The light curves of SN 2001du. The *open pentagons*, *open squares*, and *open triangles* are *BVI* measurements from images obtained using the YALO 1-m telescope (N. Suntzeff & K. Krisciunas, 2003, private communication). The *filled* symbols are the measurements from the *HST* images. Also shown for comparison are the measurements in *B* (*five-pointed stars*) and *V* (*four-pointed stars*) for the Type II-P SN 1999em from Leonard et al. (2002a), adjusted by ~ 1 mag to match the *V* peak for SN 2001du.




Fig. 4.— Hertzsprung-Russell diagram showing model stellar evolutionary tracks (alternating *long-dashed lines* and *short-dashed lines*) for a range of masses from Lejeune & Schaerer (2001), with enhanced mass loss for the most massive stars and a metallicity $Z = 0.04$. Also shown are the luminosity limits (*heavy solid lines*) for both the F555W (*V*) and F814W (*I*) bands. Stars with luminosities brighter than these limits should have been detected in the pre-SN *HST* images.

This figure "fg1.jpg" is available in "jpg" format from:

<http://arXiv.org/ps/astro-ph/0301346v2>

This figure "fg2.jpg" is available in "jpg" format from:

<http://arXiv.org/ps/astro-ph/0301346v2>